


Density functional theory (DFT)
and the concepts of the
augmented-plane-wave plus local orbital
(L)APW+lo method


Karlheinz Schwarz
Institute for Material Chemistry
TU Wien
Vienna University of Technology





DFT

Walter Kohn




APW

J.C.Slater




LAPW

O.K.Andersen




Electronic structure of solids and surfaces



hexagonal boron nitride on Rh(111)
2x2 supercell (1108 atoms per cell)

Phys.Rev.Lett. 98, 106802 (2007)



K.Schwarz, P.Blaha, S.B.Trickey,
Molecular physics, **108**, 3147 (2010)

Wien2k is used worldwide
by about 2200 groups



The WIEN2k code: comments



- **Walter Kohn:** density functional theory (DFT)
- **J.C.Slater:** augmented plane wave (APW) method, 1937
- **O.K.Andersen:** Linearized APW (LAPW)
- **Wien2k code:** developed during the last 30 years
 - *In the year 2000 (2k) the WIEN code (from Vienna) was called wien2k*
 - *One of the most accurate DFT codes for solids*
 - *All electron, relativistic, full-potential method*
 - *Widely used in academia and industry*
- **Applications:**
 - *solids: insulators , covalently bonded systems, metals*
 - *Electronic, magnetic, elastic , ...properties*
 - *Surfaces:*
 - *Many application in literature*
 - *See www.wien2k.at*



A few solid state concepts



■ Crystal structure

- *Unit cell (defined by 3 lattice vectors) leading to 7 crystal systems*
- *Bravais lattice (14)*
- *Atomic basis (Wyckoff position)*
- *Symmetries (rotations, inversion, mirror planes, glide plane, screw axis)*
- *Space group (230)*
- *Wigner-Seitz cell*
- *Reciprocal lattice (Brillouin zone)*

■ Electronic structure

- *Periodic boundary conditions*
- *Bloch theorem (k -vector), Bloch function*
- *Schrödinger equation (HF, DFT)*



Unit cell



Assuming an ideal infinite crystal we define a unit cell by


Unit cell: a volume in space that fills space entirely when translated by all lattice vectors.

The obvious choice:

a parallelepiped defined by **a**, **b**, **c**, three **basis vectors** with

the best **a**, **b**, **c** are as orthogonal as possible

the cell is as symmetric as possible (14 types)



A unit cell containing one lattice point is called **primitive cell**.



Crystal system: e.g. cubic




Axis system

$$a = b = c$$


$$\alpha = \beta = \gamma = 90^\circ$$

primitive




P (cP)

body centered



I (bcc)

face centered



F (fcc)



3D lattice types:



7 Crystal systems and 14 Bravais lattices


Triclinic	1	“no” symmetry
Monoclinic (P, C)	2	Two right angles
Orthorhombic (P, C, I, F)	4	Three right angles
Tetragonal (P, I)	2	Three right angles + 4 fold rotation
Cubic (P, I, F)	3	Three right angles + 4 fold + 3 fold
Trigonal (Rhombohedral)	1	Three equal angles ($\neq 90^\circ$) + 3 fold
Hexagonal	1	Two right and one 120° angle + 6 fold



Wigner-Seitz Cell



Form **connection** to all neighbors and **span a plane normal** to the connecting line at half distance





Bloch-Theorem:



$$\left[-\frac{1}{2} \nabla^2 + V(r) \right] \Psi(r) = E \Psi(r)$$

1-dimensional case:

$V(x)$ has lattice periodicity ("translational invariance"):

$$V(x) = V(x+a)$$

The **electron density** $\rho(x)$ has also lattice periodicity, however, the **wave function** does **NOT**:

$$\rho(x) = \rho(x+a) = \Psi^*(x)\Psi(x) \quad \text{but:}$$

$$\Psi(x+a) = \mu \Psi(x) \quad \Rightarrow \quad \mu^* \mu = 1$$


Application of the translation τ g -times:

$$\tau^g \Psi(x) = \Psi(x+ga) = \mu^g \Psi(x)$$



- The wave function must be uniquely defined: after G translations it must be identical ($G a$: periodicity volume):

$$\tau^G \Psi(x) = \Psi(x + Ga) = \mu^G \Psi(x) = \Psi(x)$$
$$\Rightarrow \mu^G = 1$$



$$\mu = e^{2\pi i \frac{g}{G}} \quad g = 0, \pm 1, \pm 2, \dots$$

$$\text{Def.} : \quad k = \frac{2\pi}{a} \frac{g}{G} \quad \mu = e^{ika}$$

$$\text{Bloch condition} : \quad \Psi(x + a) = e^{ika} \Psi(x) = \Psi_k$$



Bloch functions:




Wave functions with Bloch form:

$$\Psi_k(x) = e^{ikx} u(x) \quad \text{where:} \quad u(x) = u(x+a)$$

Phase factor

lattice periodic function



Replacing k by $k+K$, where K is a **reciprocal lattice vector**, fulfills again the Bloch-condition.


→ k can be restricted to the **first Brillouin zone**.


$$e^{i\frac{2\pi}{a}K} = 1$$

$$-\frac{\pi}{a} < k < \frac{\pi}{a}$$



Concepts when solving Schrödingers-equation in solids







DFT vs. MBT (many body theory)



Two communities in solid state theory


	LDA bandstructure	many body theory
+	<ul style="list-style-type: none">material-specific, "ab initio"often successful, quantitative	<ul style="list-style-type: none">electronic correlationsqualitative understanding
-	<ul style="list-style-type: none">effective one-particle approach	<ul style="list-style-type: none">model Hamiltonian

time averaged
electron density



⤴: Coulomb WW
⇒ correlations

←
lattice pot.



Ab-initio Hamiltonian

(non-relativistic/Born-Oppenheimer approximation)



kinetic energy


lattice potential

Coulomb interaction

$$H = \sum_i \left[-\frac{\hbar^2 \Delta_i}{2m_e} + \sum_l \frac{-e^2}{4\pi\epsilon_0} \frac{Z_l}{|\mathbf{r}_i - \mathbf{R}_l|} \right] + \frac{1}{2} \sum_{i \neq j} \frac{e^2}{4\pi\epsilon_0} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$$

LDA bandstructure corresponds to

$$H_{\text{LDA}} = \sum_i \left[-\frac{\hbar^2 \Delta_i}{2m_e} + \sum_l \frac{-e^2}{4\pi\epsilon_0} \frac{1}{|\mathbf{r}_i - \mathbf{R}_l|} + \int d^3r \frac{e^2}{4\pi\epsilon_0} \frac{1}{|\mathbf{r}_i - \mathbf{r}|} \rho(\mathbf{r}) + V_{xc}^{\text{LDA}}(\rho(\mathbf{r}_i)) \right]$$



Coulomb potential:

- nuclei
- all electrons
- including self-interaction

Quantum mechanics:


- exchange
- correlation
- (partly) cancel self-interaction



ESSENCE OF DENSITY-FUNCTIONAL THEORY



- **Every observable quantity of a quantum system can be calculated from the density of the system ALONE (Hohenberg, Kohn, 1964).**
- **The density of particles interacting with each other can be calculated as the density of an auxiliary system of non-interacting particles (Kohn, Sham, 1965).**







Walter Kohn, Nobel Prize 1998 Chemistry



Walter Kohn



**“Self-consistent Equations including Exchange and Correlation Effects”
W. Kohn and L. J. Sham, Phys. Rev. 140, A1133 (1965)**

Literal quote from Kohn and Sham’s paper: “... We do not expect an accurate description of chemical binding.”



Hohenberg-Kohn theorem: (exact)

The total energy of an interacting inhomogeneous electron gas in the presence of an external potential $V_{\text{ext}}(\mathbf{r})$ is a **functional** of the density ρ

$$E = \int V_{\text{ext}}(\mathbf{r}) \rho(\mathbf{r}) d\mathbf{r} + F[\rho]$$

Kohn-Sham: (still exact!)

$$E = T_o[\rho] + \int V_{\text{ext}} \rho(\mathbf{r}) d\mathbf{r} + \frac{1}{2} \int \frac{\rho(\mathbf{r}) \rho(\mathbf{r}')}{|\mathbf{r}' - \mathbf{r}|} d\mathbf{r} d\mathbf{r}' + E_{\text{xc}}[\rho]$$

E_{kinetic}
non interacting

E_{ne}

E_{coulomb} E_{ee}


E_{xc} exchange-correlation

In KS the many body problem of interacting electrons and nuclei is mapped to a one-electron reference system that leads to the same density as the real system.



Exchange and correlation

- We divide the density of the N-1 electron system into the total density $n(r)$ and an exchange-correlation hole:



Properties of the exchange-correlation hole:

- Locality
 - Pauli principle
 - the hole contains ONE electron
 - The hole must be negative
-
- The exchange hole affects electrons with the same spin and accounts for the Pauli principle
 - In contrast, the correlation-hole accounts for the Coulomb repulsion of electrons with the opposite spin. It is short range and leads to a small redistribution of charge. The correlation hole contains NO charge:

$$h(r_0, r) \xrightarrow{|r-r_0| \rightarrow \infty} 0$$

$$h(r_0, r) \xrightarrow{|r-r_0| \rightarrow 0} -n(r_0)$$

$$\int dr h(r_0, r) = -1$$

$$h(r_0, r) \leq 0$$

$$\int dr h_c(r_0, r) = 0$$



Kohn-Sham equations



LDA, GGA

$$E = T_o[\rho] + \int V_{ext} \rho(\vec{r}) d\vec{r} + \frac{1}{2} \int \frac{\rho(\vec{r})\rho(\vec{r}')}{|\vec{r}' - \vec{r}|} d\vec{r} d\vec{r}' + E_{xc}[\rho]$$

1-electron equations (Kohn Sham)

vary ρ

$$\left\{ -\frac{1}{2} \nabla^2 + V_{ext}(\vec{r}) + V_C(\rho(\vec{r})) + V_{xc}(\rho(\vec{r})) \right\} \Phi_i(\vec{r}) = \varepsilon_i \Phi_i(\vec{r})$$

$$-Z/r$$

$$\int \frac{\rho(\vec{r}')}{|\vec{r}' - \vec{r}|} d\vec{r}'$$

$$\frac{\partial E_{xc}(\rho)}{\partial \rho}$$

$$\rho(\vec{r}) = \sum_{\varepsilon_i \leq E_F} |\Phi_i|^2$$

$$E_{xc}^{LDA} \propto \int \rho(r) \varepsilon_{xc}^{hom.}[\rho(r)] dr$$

$$E_{xc}^{GGA} \propto \int \rho(r) F[\rho(r), \nabla \rho(r)] dr$$

LDA


GGA

treats both,
exchange and correlation effects,
but approximately

New (better ?) functionals are still an active field of research



DFT ground state of iron



LSDA

- *NM*
- *fcc*
- *in contrast to experiment*

GGA

- *FM*
- *bcc*
- *Correct lattice constant*

Experiment

- *FM*
- *bcc*




DFT thanks to Claudia Ambrosch (Graz)




GGA follows LDA







CoO AFM-II total energy, DOS



■ CoO

- *in NaCl structure*
- *antiferromagnetic: AF II*
- *insulator*
- *t_{2g} splits into a_{1g} and e_g'*
- *GGA almost splits the bands*






CoO why is GGA better than LSDA




$$\Delta V_{xc}^{\uparrow} = V_{xc}^{\uparrow GGA} - V_{xc}^{\uparrow LSDA}$$




■ Central Co atom distinguishes


- *between* Co^{\uparrow}
- *and* Co^{\downarrow}

■ Angular correlation







FeF₂: GGA works surprisingly well



LSDA



GGA



Fe-EFG in FeF₂:

LSDA: 6.2

GGA: 16.8

exp: 16.5

FeF₂: GGA splits
t_{2g} into a_{1g} and e_g'

agree



Accuracy of DFT for transition metals



Lattice parameters (Å)

	Exp.	LDA	PBE	WC
Co	2.51	2.42	2.49	2.45
Ni	3.52	3.42	3.52	3.47
Cu	3.61	3.52	3.63	3.57
Ru	2.71	2.69	2.71	2.73
Rh	3.80	3.76	3.83	3.80
Pd	3.88	3.85	3.95	3.89
Ag	4.07	4.01	4.15	4.07
Ir	3.84	3.84	3.90	3.86
Pt	3.92	3.92	4.00	3.96
Au	4.08	4.07	4.18	4.11

■ 3d elements:

- *PBE superior, LDA much too small*

■ 4d elements:

LDA too small, PBE too large

- *New functional Wu-Cohen (WC)*

Z.Wu, R.E.Cohen,
PRB 73, 235116 (2006)

■ 5d elements:

- *LDA superior, PBE too large*




accuracy: "DFT limit"



- Testing of DFT functionals:
 - error of theoretical lattice parameters for a large variety of solids (Li-Th)

	me (Å)	mae (Å)	mre (%)	mare (%)
LDA	-0.058	0.058	-1.32	1.32
SO-GGA	-0.014	0.029	-0.37	0.68
PBEsol	-0.005	0.029	-0.17	0.67
WC	0.000	0.031	-0.03	0.68
AM05	0.005	0.035	0.01	0.77
PBE	0.051	0.055	1.05	1.18






Can LDA be improved ?



- **better GGAs and meta-GGAs ($\rho, \nabla\rho, \tau$):**
 - *usually improvement, but often too small.*
- **LDA+U:** for correlated 3d/4f electrons, treat strong Coulomb repulsion via Hubbard U parameter (cheap, “empirical U” ?)
- **Exact exchange:** imbalance between exact X and approximate C
 - *hybrid-DFT (mixing of HF + GGA; “mixing factor” ?)*
 - *exact exchange + RPA correlation (extremely expensive)*
- **GW:** gaps in semiconductors, expensive!
- **Quantum Monte-Carlo:** very expensive
- **DMFT:** for strongly correlated (metallic) d (f) -systems (expensive)



Application to FeO



Table: Lattice constant a (Å), bulk modulus B (GPa), total and orbital magnetic moment M and M_l (μ_B), fundamental band gap Δ_{fund} (eV), and optical band gap Δ_{opt} (eV) of AFII phase of FeO.

	a	B	M (M_l)	Δ_{fund}	Δ_{opt}	
LDA	4.18	230	3.44 (0.09)	0.0	0.0	} metallic
PBE	4.30	183	3.49 (0.08)	0.0	0.0	
LDA+ U	4.28	199	4.23 (0.63)	1.7	2.2	} gap
B3PW91	4.35	172	4.15 (0.61)	1.3	1.8	
PBE0	4.40	155	4.30 (0.75)	1.2	1.6	
Fock-0.35	4.31	195	4.27 (0.68)	2.1	2.4	
Fock-0.5	4.34	189	4.32 (0.68)	2.2	2.7	
Expt.	4.334	150–180	3.32, 4.2	2.4	0.5 ¹ , 2.4 ²	


¹Assigned to Fe 3d/O 2sp → Fe 4s transitions.

²Assigned to Fe 3d/O 2sp → Fe 3d transitions.

F. Tran, P. Blaha, K. Schwarz, P. Novák, PRB **74**, 155108 (2006)



FeO: LDA vs. LDA+U vs. Hybrids vs. exp





Band gaps by a semi-local potential



- **Becke-Johnson potential** (J. Chem. Phys. 124, 221101 (2006))
 - *cheap* potential designed to reproduce expensive exact X-OEP potentials in atoms
 - tests for bandgaps in solids (Tran, Blaha, Schwarz, J. Phys. CM. 19, 196208 (2007))
gives only small improvement over LDA/GGA gaps
- **modified Becke-Johnson potential**

$$v_{x,\sigma}^{\text{MBJ}}(\mathbf{r}) = c v_{x,\sigma}^{\text{BR}}(\mathbf{r}) + (3c - 2) \frac{1}{\pi} \sqrt{\frac{5}{12}} \sqrt{\frac{2t_{\sigma}(\mathbf{r})}{\rho_{\sigma}(\mathbf{r})}}$$

$$t_{\sigma} = (1/2) \sum_{i=1}^{N_{\sigma}} \nabla \psi_{i,\sigma}^* \cdot \nabla \psi_{i,\sigma}$$

$$v_{x,\sigma}^{\text{BR}}(\mathbf{r}) = -\frac{1}{b_{\sigma}(\mathbf{r})} \left(1 - e^{-x_{\sigma}(\mathbf{r})} - \frac{1}{2} x_{\sigma}(\mathbf{r}) e^{-x_{\sigma}(\mathbf{r})} \right)$$

$$c = \alpha + \beta \left(\frac{1}{V_{\text{cell}}} \int_{\text{cell}} \frac{|\nabla \rho(\mathbf{r}')|}{\rho(\mathbf{r}')} d^3 r' \right)^{1/2}$$

F. Tran P. Blaha
PRL 102, 226401 (2009)

kinetic energy density


Becke-Roussel potential
(approximate Slater-pot.)

c depends on the density
properties of a material



band gaps by mBJ-LDA

- LDA strongly underestimates gaps
- mBJ gives gaps of „GW quality“





Treatment of exchange and correlation



Approximations for E_{xc}

- ▶ LDA: $E_{xc}^{LDA} = \int f(\rho(\mathbf{r}))d^3r$
- ▶ GGA: $E_{xc}^{GGA} = \int f(\rho(\mathbf{r}), |\nabla\rho(\mathbf{r})|)d^3r$
- ▶ MGGA: $E_{xc}^{MGGA} = \int f(\rho(\mathbf{r}), |\nabla\rho(\mathbf{r})|, \nabla^2\rho(\mathbf{r}), t(\mathbf{r}))d^3r$
- ▶ LDA+U: $E_{xc}^{LDA+U} = E_{xc}^{LDA} + E_{ee} - E_{dc}$
- ▶ GGA+U: $E_{xc}^{GGA+U} = E_{xc}^{GGA} + E_{ee} - E_{dc}$
- ▶ hybrid: $E_{xc}^{hybrid} = E_{xc}^{DFT} + \alpha (E_x^{HF} - E_x^{DFT})$
where

$$E_x^{HF} = -\frac{1}{2} \sum_{\sigma} \sum_{\substack{n,k \\ n',k'}} w_k w_{k'} \int \int \frac{\psi_{nk}^{\sigma*}(\mathbf{r}) \psi_{n'k'}^{\sigma*}(\mathbf{r}') \psi_{n'k'}^{\sigma}(\mathbf{r}) \psi_{nk}^{\sigma}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d^3r d^3r'$$





Hybrid functional: only for (correlated) electrons



- Only for certain atoms and electrons of a given angular momentum ℓ

$$E_{xc}^{\text{hybrid}} = E_{xc}^{\text{DFT}}[\rho^\sigma] + \alpha \left(E_x^{\text{HF}}[n_{m_i m_j}^\sigma] - E_x^{\text{DFT}}[\rho_\ell^\sigma] \right)$$

$$E_x^{\text{HF}}[n_{m_i m_j}^\sigma] = -\frac{1}{2} \sum_{\sigma} \sum_{m_1, m_2, m_3, m_4}^{\ell} n_{m_1 m_2}^\sigma n_{m_3 m_4}^\sigma \langle m_1 m_3 | v_{ee} | m_4 m_2 \rangle$$



$$\langle m_1 m_2 | v_{ee} | m_3 m_4 \rangle = \sum_{k=0}^{2\ell} a_k F_k$$

The Slater integrals F_k are calculated according to P. Novák et al., phys.stat.sol (b) **245**, 563 (2006)



Structure: $a, b, c, \alpha, \beta, \gamma, R_\alpha, \dots$

unit cell atomic positions



$\mathbf{k} \in \text{IBZ}$ (irred. Brillouin zone)

Kohn Sham

$$[-\nabla^2 + V(\rho)]\psi_k = E_k \psi_k$$

$$\psi_k = \sum_{k_n} C_{k_n} \Phi_{k_n}$$

Variational method

$$\frac{\delta \langle E \rangle}{\delta C_{k_n}} = 0$$

Generalized eigenvalue problem

$$HC = ESC$$

$$\rho = \sum_{E_k \leq E_F} \psi_k^* \psi_k$$



Solving Schrödinger's equation:

$$\left[-\frac{1}{2} \nabla^2 + V(r) \right] \Psi_i^k = \varepsilon_i^k \Psi_i^k$$



- Ψ cannot be found analytically
- complete "numerical" solution is possible but inefficient

■ Ansatz:

- *linear combination of some "basis functions"*
 - different methods use different basis sets !
- *finding the "best" wave function using the **variational** principle:*

$$\Psi_k = \sum_{K_n} c_{k_n} \Phi_{k_n}$$

$$\langle E_k \rangle = \frac{\langle \Psi_k^* | H | \Psi_k \rangle}{\langle \Psi_k^* | \Psi_k \rangle} \quad \frac{\partial E_k}{\partial c_{k_n}} = 0$$

- *this leads to the famous "Secular equations", i.e. a set of linear equations which in matrix representation is called "generalized eigenvalue problem"*

$$H C = E S C$$

H, S : hamilton and overlap matrix; C: eigenvectors, E: eigenvalues




■ plane waves

- *pseudo potentials*
- *PAW (projector augmented wave) by P.E.Blöchl*


■ space partitioning (augmentation) methods

- *LMTO (linear muffin tin orbitals)*
 - ASA approx., linearized numerical radial function + Hankel- and Bessel function expansions
 - full-potential LMTO
- *ASW (augmented spherical wave)*
 - similar to LMTO
- *KKR (Korringa, Kohn, Rostocker method)*
 - solution of multiple scattering problem, Greens function formalism
 - equivalent to APW
- *(L)APW (linearized augmented plane waves)*



■ LCAO methods

- *Gaussians, Slater, or numerical orbitals, often with PP option)*






pseudopotential plane wave methods



- **plane waves** form a “complete” basis set, however, they “never” converge due to the rapid oscillations of the atomic wave functions χ close to the nuclei
- let’s get rid of all **core electrons** and **these oscillations** by replacing the strong ion–electron potential by a much weaker (and physically dubious) ***pseudopotential***
- **Hellmann’s** 1935 *combined approximation method*







"real" potentials vs. pseudopotentials



- "real" potentials contain the **Coulomb singularity** $-Z/r$
- the wave function has a **cusp** and many **wiggles**,
- **chemical bonding** depends mainly on the overlap of the wave functions between neighboring atoms (in the region between the nuclei) \rightarrow





APW based schemes



■ APW (J.C.Slater 1937)

- *Non-linear eigenvalue problem*
- *Computationally very demanding*

K.Schwarz, P.Blaha, G.K.H.Madsen,
Comp.Phys.Commun. **147**, 71-76 (2002)

■ LAPW (O.K.Anderssen 1975)

- *Generalized eigenvalue problem*
- *Full-potential*


K.Schwarz,
DFT calculations of solids with LAPW and WIEN2k
Solid State Chem. **176**, 319-328 (2003)

■ Local orbitals (D.J.Singh 1991)


- *treatment of semi-core states (avoids ghostbands)*

■ APW+lo (E.Sjöstedt, L.Nordström, D.J.Singh 2000)

- *Efficiency of APW + convenience of LAPW*
- *Basis for*



K.Schwarz, P.Blaha, S.B.Trickey,
Molecular physics, **108**, 3147 (2010)





APW Augmented Plane Wave method



The unit cell is partitioned into:

atomic spheres
Interstitial region



Basis set:

PW: $e^{i(\vec{k} + \vec{K}) \cdot \vec{r}}$

Atomic partial waves


$$\sum_{lm} A_{lm}^K u_l(r', \varepsilon) Y_{lm}(\hat{r}')$$

join

$u_l(r, \varepsilon)$ are the numerical solutions of the radial Schrödinger equation in a given spherical potential for a particular energy ε
 A_{lm}^K coefficients for matching the PW



Slater's APW (1937)



H Hamiltonian
S overlap matrix


Atomic partial waves

$$\sum_{\ell m} a_{\ell m}^K u_{\ell}(r', \varepsilon) Y_{\ell m}(\hat{r}')$$

Energy dependent basis functions
lead to a

Non-linear eigenvalue problem


Numerical search for those energies, for which the $\det|H-ES|$ vanishes. **Computationally very demanding.**
"Exact" solution for given MT potential!



Linearization of energy dependence

LAPW suggested by

O.K.Andersen,
Phys.Rev. B 12, 3060
(1975)




$$\Phi_{k_n} = \sum_{\ell m} [A_{\ell m}(k_n) u_{\ell}(E_{\ell}, r) + B_{\ell m}(k_n) \dot{u}_{\ell}(E_{\ell}, r)] Y_{\ell m}(\hat{r})$$

expand u_l at fixed energy E_l and
add $\dot{u}_l = \partial u_l / \partial \epsilon$

$A_{\ell m}^k, B_{\ell m}^k$: join PWs in
value and slope

→ General eigenvalue problem
(diagonalization)

→ additional constraint requires
more PWs than APW






shape approximations to "real" potentials




■ Atomic sphere approximation (ASA)

- *overlapping spheres "fill" all volume*
- *potential spherically symmetric*




■ "muffin-tin" approximation (MTA)

- *non-overlapping spheres with spherically symmetric potential +*
- *interstitial region with $V=const.$*




■ "full"-potential


- *no shape approximations to V*




Full-potential in LAPW (A.Freeman et al)



SrTiO₃



Full potential



Muffin tin approximation





TiO₂ rutile

- The potential (and charge density) can be of general form (no shape approximation)

$$V(r) = \begin{cases} \sum_{LM} V_{LM}(r) Y_{LM}(\hat{r}) & r < R_\alpha \\ \sum_K V_K e^{i\vec{K}\cdot\vec{r}} & r \in I \end{cases}$$

- Inside each atomic sphere a local coordinate system is used (defining LM)






Core, semi-core and valence states




For example: **Ti**



1s, 2s, 2p
-356.6 -38.3 -31.7 Ry

1 Ry = 13.605 eV


- **Valences states**
 - *High in energy*
 - *Delocalized* wavefunctions
- **Semi-core states**
 - *Medium energy*
 - *Principal **QN** one less than valence (e.g. in Ti **3p** and **4p**)*
 - *not completely confined inside sphere (charge leakage)*
- **Core states**
 - *Low in energy*
 - *Reside inside sphere*



Local orbitals (LO)



Ti atomic sphere



$$\Phi_{LO} = [A_{lm}u_l^{E_1} + B_{lm}\dot{u}_l^{E_1} + C_{lm}u_l^{E_2}]Y_{lm}(\hat{r})$$

■ LOs

- *are confined to an atomic sphere*
- *have zero value and slope at R*
- *Can treat two principal QN n for each azimuthal QN ℓ (e.g. 3p and 4p)*
- *Corresponding states are strictly orthogonal*
 - (e.g. semi-core and valence)
- *Tail of semi-core states can be represented by plane waves*
- *Only slightly increases the basis set (matrix size)*

D.J.Singh,
Phys.Rev. B 43 6388 (1991)



An alternative combination of schemes



E.Sjöstedt, L.Nordström, D.J.Singh,
An alternative way of linearizing the augmented plane wave method,
Solid State Commun. 114, 15 (2000)

- Use **APW**, but at **fixed E_l** (superior PW convergence)
- **Linearize** with **additional local orbitals (lo)**
(add a few extra basis functions)

$$\Phi_{k_n} = \sum_{lm} A_{lm}(k_n) u_l(E_l, r) Y_{lm}(\hat{r})$$

$$\Phi_{lo} = [A_{lm} u_l^{E_1} + B_{lm} \dot{u}_l^{E_1}] Y_{lm}(\hat{r})$$

optimal solution: mixed basis


- use APW+lo for states, which are difficult to converge:
(f or d- states, atoms with small spheres)
- use LAPW+LO for all other atoms and angular momenta



Improved convergence of APW+lo



Representative Convergence:




e.g. force (F_y) on oxygen in SES vs. # plane waves:

- in **LAPW** changes sign and converges slowly
- in **APW+lo** better convergence
- to same value as in LAPW

SES (sodium electro solodalite)

K.Schwarz, P.Blaha, G.K.H.Madsen,
Comp.Phys.Commun. **147**, 71-76 (2002)





Summary: Linearization LAPW vs. APW



Atomic partial waves

LAPW

$$\Phi_{k_n} = \sum_{\ell m} [A_{\ell m}(k_n)u_{\ell}(E_{\ell}, r) + B_{\ell m}(k_n)\dot{u}_{\ell}(E_{\ell}, r)]Y_{\ell m}(\hat{r})$$

APW+lo

$$\Phi_{k_n} = \sum_{\ell m} A_{\ell m}(k_n)u_{\ell}(E_{\ell}, r)Y_{\ell m}(\hat{r})$$

plus another type of local orbital (lo)

Plane Waves (PWs)

$$e^{i(\vec{k} + \vec{K}_n) \cdot \vec{r}}$$


match at sphere boundary

LAPW

value and slope $A_{\ell m}(k_n), B_{\ell m}(k_n)$

APW

value $A_{\ell m}(k_n)$





E.Sjöststedt, L.Nordström, D.J.Singh, SSC 114, 15 (2000)

- Use **APW**, but at **fixed E_f** (superior PW convergence)
- **Linearize** with **additional lo** (add a few basis functions)

optimal solution: mixed basis


- use **APW+lo** for states which are difficult to converge:
(**f-** or **d-** states, atoms with small spheres)
- use **LAPW+LO** for all **other** atoms and angular momenta

A summary is given in

K.Schwarz, P.Blaha, G.K.H.Madsen,
Comp.Phys.Commun. **147**, 71-76 (2002)



The WIEN2k authors



**An Augmented Plane Wave
Plus Local Orbital Program for
Calculating Crystal Properties**

**Peter Blaha
Karlheinz Schwarz
Georg Madsen
Dieter Kvasnicka
Joachim Luitz**

November 2001
Vienna, AUSTRIA
Vienna University of Technology



G.Madsen

P.Blaha

D.Kvasnicka

K.Schwarz

J.Luitz

<http://www.wien2k.at>




International users

about **2200** licenses worldwide

Europe: A, B, CH, CZ, D, DK, ES, F, FIN, GR, H, I, IL, IRE, N, NL, PL, RO, S, SK, SL, SI, UK (ETH Zürich, MPI Stuttgart, FHI Berlin, DESY, RWTH Aachen, ESRF, Prague, IJS Ljubljana, Paris, Chalmers, Cambridge, Oxford)

America: ARG, BZ, CDN, MX, USA (MIT, NIST, Berkeley, Princeton, Harvard, Argonne NL, Los Alamos NL, Oak Ridge NL, Penn State, Purdue, Georgia Tech, Lehigh, John Hopkins, Chicago, Stony Brook, SUNY, UC St.Barbara, UCLA)

far east: AUS, China, India, JPN, Korea, Pakistan, Singapore, Taiwan (Beijing, Tokyo, Osaka, Kyoto, Sendai, Tsukuba, Hong Kong)



75 industries (Canon, Eastman, Exxon, Fuji, Hitachi, IBM, Idemitsu Petrochem., Kansai, Komatsu, Konica-Minolta, A.D.Little, Mitsubishi, Mitsui Mining, Motorola, NEC, Nippon Steel, Norsk Hydro, Osram, Panasonic, Samsung, Seiko Epson, Siemens, Sony, Sumitomo, TDK, Toyota).

mailinglist: 10.000 emails/6 years



The first publication of the WIEN code



FULL-POTENTIAL, LINEARIZED AUGMENTED PLANE WAVE PROGRAMS FOR CRYSTALLINE SYSTEMS

P. BLAHA, K. SCHWARZ, and P. SORANTIN

Institut für Technische Elektrochemie, Technische Universität Wien, A-1060 WIEN, Austria

and


S.B. TRICKEY

Quantum Theory Project, Depts. of Physics and of Chemistry, University of Florida, Gainesville, FL 32611, USA

PROGRAM SUMMARY

Title of program: WIEN

Computer Physics Communications 59 (1990) 399–415






Europa Austria Vienna → WIEN



In the Heart of EUROPE






In Japan

- Book published by
Shinya Wakoh (2006)

『WIEN2k 入門』追加版

改訂 固体の中の電子

バンド計算の基礎と応用



和光システム研究所 著

WIEN2k 入門

WIEN-code は 1980 年ごろから、グループの指導者である Karlheinz Schwarz によって書き始められ、1990 年に最初の copyrighted version の WIEN が発表された。その後 UNIX version となり、WIEN93, WIEN95, WIEN97 を経て、Fortran90 対応の WIEN2k へと改良・拡張されてきた*1。基礎となるシュレーディンガー方程式はコーン・シャム方程式であり、バンド計算法は主として FLAPW 法、ポテンシャルは LSDA, GGA などである。最新の WIEN2k では、APW+lo も取り入れられており、ポテンシャルとしては電子相関が強いときに必要であると云われている補正 +U も扱えるようになっている。また、並列計算機を使えば、極めて複雑な結晶も計算の対象とすることができる。



Development of WIEN2k




■ Authors of WIEN2k

P. Blaha, K. Schwarz, D. Kvasnicka, G. Madsen and J. Luitz

■ Other contributions to WIEN2k

- *C. Ambrosch-Draxl (Univ. Graz, Austria), optics*
- *T. Charpin (Paris), elastic constants*
- *R. Laskowski (Vienna), non-collinear magnetism, parallelization*
- *L. Marks (Northwestern, US), various optimizations, new mixer*
- *P. Novák and J. Kunes (Prague), LDA+U, SO*
- *B. Olejnik (Vienna), non-linear optics,*
- *C. Persson (Uppsala), irreducible representations*
- *V. Petricek (Prague) 230 space groups*
- *M. Scheffler (Fritz Haber Inst., Berlin), forces*
- *D.J.Singh (NRL, Washington D.C.), local orbitals (LO), APW+lo*
- *E. Sjöstedt and L Nordström (Uppsala, Sweden), APW+lo*
- *J. Sofo and J. Fuhr (Barriloche), Bader analysis*
- *B. Yanchitsky and A. Timoshevskii (Kiev), spacegroup*



■ and many others



A series of WIEN workshops were held



■ 1st	Vienna	April	1995	Wien95
■ 2nd	Vienna	April	1996	
■ 3rd	Vienna	April	1997	Wien97
■ 4st	Trieste, Italy	June	1998	
■ 5st	Vienna	April	1999	
■ 6th	Vienna	April	2000	
■ 7th	Vienna	Sept.	2001	Wien2k
■ 8th	Esfahan, Iran	April	2002	
■	Penn State, USA	July	2002	
■ 9th	Vienna	April	2003	
■ 10th	Penn State, USA	July	2004	
■ 11th	Kyoto, Japan	May	2005	
■	IPAM, Los Angeles, USA	Nov.	2005	
■ 12th	Vienna	April	2006	
■ 13th	Penn State, USA	June	2007	
■ 14th	Singapore	July	2007	
■ 15th	Vienna	March	2008	
■ 16th	Penn State, USA	June	2009	
■ 17th	Nantes, France	July	2010	
■ 18th	Penn State, USA	June	2011	
■ 19th	Tokyo, Japan	Sept	2012	
■ 20th	Penn State, USA	Aug.	2013	





APW + local orbital method
(linearized) augmented plane wave method

Total wave function $\Psi_k = \sum_{K_n} C_{k_n} \phi_{k_n}$ n...50-100 PWs /atom

Variational method:

$$\langle E \rangle = \frac{\langle \Psi | H | \Psi \rangle}{\langle \Psi | \Psi \rangle} \quad \frac{\delta \langle E \rangle}{\delta C_{k_n}} = 0$$

upper bound

minimum

Generalized eigenvalue problem:



$$H C = E S C$$

Diagonalization of (real or complex) matrices of size 10.000 to 50.000 (up to 50 Gb memory)



Structure: $a, b, c, \alpha, \beta, \gamma, R_\alpha, \dots$

unit cell atomic positions



$\mathbf{k} \in \text{IBZ}$ (irred. Brillouin zone)

Kohn Sham

$$[-\nabla^2 + V(\rho)]\psi_k = E_k \psi_k$$

$$\psi_k = \sum_{k_n} C_{k_n} \Phi_{k_n}$$

Variational method

$$\frac{\delta \langle E \rangle}{\delta C_{k_n}} = 0$$

Generalized eigenvalue problem

$$HC = ESC$$

$$\rho = \sum_{E_k \leq E_F} \psi_k^* \psi_k$$



The Brillouin zone (BZ)




■ Irreducible BZ (IBZ)

- *The irreducible wedge*
- *Region, from which the whole BZ can be obtained by applying all symmetry operations*

■ Bilbao Crystallographic Server:

- www.cryst.ehu.es/cryst/
- *The IBZ of all space groups can be obtained from this server*
- *using the option KVEC and specifying the space group (e.g. No.225 for the fcc structure leading to bcc in reciprocal space, No.229)*







Self-consistent field (SCF) calculations




- In order to solve $H\Psi = E\Psi$ we need to know the potential $V(r)$
- for $V(r)$ we need the electron density $\rho(r)$
- the density $\rho(r)$ can be obtained from $\Psi(r)^*\Psi(r)$
- ?? $\Psi(r)$ is unknown before $H\Psi = E\Psi$ is solved ??






Band structure of fcc Cu





Flow Chart of WIEN2k (SCF)




WIEN2k: *P. Blaha, K. Schwarz, G. Madsen, D. Kvasnicka, and J. Luitz*



Workflow of a WIEN2k calculation



- individual FORTRAN programs linked by shell-scripts
- the output of one program is input for the next
- lapw1/2 can run in parallel on many processors




* fraction of total computation time



Advantage/disadvantage of WIEN2k




- + robust all-electron full-potential method (new effective mixer)
- + unbiased basis set, one convergence parameter (LDA-limit)
- + all elements of periodic table (comparable in CPU time), metals
- + LDA, GGA, meta-GGA, LDA+U, spin-orbit
- + many properties and tools (supercells, symmetry)
- + w2web (for novice users)
- ? speed + memory requirements
 - + *very efficient basis for large spheres (2 bohr) (Fe: 12Ry, O: 9Ry)*
 - *less efficient for small spheres (1 bohr) (O: 25 Ry)*
 - *large cells, many atoms (n^3 , but new iterative diagonalization)*
 - *full H, S matrix stored \rightarrow large memory required*
 - + *effective dual parallelization (k-points, mpi-fine-grain)*
 - + *many k-points do not require more memory*
- no stress tensor
- no linear response



w2web GUI (graphical user interface)



- **Structure generator**
 - *spacegroup selection*
 - *import cif file*
- **step by step initialization**
 - *symmetry detection*
 - *automatic input generation*
- **SCF calculations**
 - *Magnetism (spin-polarization)*
 - *Spin-orbit coupling*
 - *Forces (automatic geometry optimization)*
- **Guided Tasks**
 - *Energy band structure*
 - *DOS*
 - *Electron density*
 - *X-ray spectra*
 - *Optics*



Execution >>

StructGen™
initialize calc.
run SCF
single prog.
optimize(V,c/a)
mini. positions

Utils. >>

Tasks >>

Files >>

struct file(s)
input files
output files
SCF files

Session Mgmt. >>

change session
change dir
change info

Configuration

Usersguide

html-Version
pdf-Version

Idea and realization
by

Session: TiC

/area51/pblaha/lapw/2005-june/TiC

StructGen™

You have to click "Save Structure" for changes to take effect!

Save Structure

Title: TiC

Lattice:

Type: F

P
F
B
CXY
CYZ
CXZ
R
H
1_P1

Spacegroups from
Bilbao Cryst Server

Lattice parameters in Å

a=4.328000038 b=4.328000038 c=4.328000038

α =90.000000 β =90.000000 γ =90.000000

Inequivalent Atoms: 2

Atom 1: Ti Z=22.0 RMT=2.0000 remove atom

Pos 1: x=0.00000000 y=0.00000000 z=0.00000000 remove
add position

Atom 2: C Z=6.0 RMT=1.9000 remove atom

Pos 1: x=0.50000000 y=0.50000000 z=0.50000000 remove
add position




Spacegroup $P4_2/mnm$

Structure given by:
 spacegroup
 lattice parameter
 positions of atoms
 (basis)

Rutile TiO_2 :
 $P4_2/mnm$ (136)
 $a=8.68, c=5.59$ bohr
Ti: (0,0,0)

O: (0.304,0.304,0)
 Wyckoff position: $x, x, 0$




$P4_2/mnm$
 D_{4h}^{14}

No. 136

$P 4_2/m 2_1/n 2/m$

$4/m m m$ Tetragonal




Origin at centre (mmm)

Number of positions,
 Wyckoff notation,
 and point symmetry

Co-ordinates of equivalent positions


Conditions limiting
 possible reflections

Number of positions, Wyckoff notation, and point symmetry			Co-ordinates of equivalent positions	Conditions limiting possible reflections
16	k	1	$x, y, z; \bar{x}, \bar{y}, z; \frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} + z; \frac{1}{2} - x, \frac{1}{2} + y, \frac{1}{2} + z;$ $x, y, \bar{z}; \bar{x}, \bar{y}, \bar{z}; \frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} - z; \frac{1}{2} - x, \frac{1}{2} + y, \frac{1}{2} - z;$ $y, x, z; \bar{y}, \bar{x}, z; \frac{1}{2} + y, \frac{1}{2} - x, \frac{1}{2} + z; \frac{1}{2} - y, \frac{1}{2} + x, \frac{1}{2} + z;$ $y, x, \bar{z}; \bar{y}, \bar{x}, \bar{z}; \frac{1}{2} + y, \frac{1}{2} - x, \frac{1}{2} - z; \frac{1}{2} - y, \frac{1}{2} + x, \frac{1}{2} - z.$	General: hkl : No conditions $hk0$: No conditions $0kl$: $k+l=2n$ hhl : No conditions
8	j	m	$x, x, z; \bar{x}, \bar{x}, z; \frac{1}{2} + x, \frac{1}{2} - x, \frac{1}{2} + z; \frac{1}{2} - x, \frac{1}{2} + x, \frac{1}{2} + z;$ $x, x, \bar{z}; \bar{x}, \bar{x}, \bar{z}; \frac{1}{2} + x, \frac{1}{2} - x, \frac{1}{2} - z; \frac{1}{2} - x, \frac{1}{2} + x, \frac{1}{2} - z.$	Special: as above, plus } no extra conditions
8	i	m	$x, y, 0; \bar{x}, \bar{y}, 0; \frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2}; \frac{1}{2} - x, \frac{1}{2} + y, \frac{1}{2};$ $y, x, 0; \bar{y}, \bar{x}, 0; \frac{1}{2} + y, \frac{1}{2} - x, \frac{1}{2}; \frac{1}{2} - y, \frac{1}{2} + x, \frac{1}{2}.$	
8	h	2	$0, \frac{1}{2}, z; 0, \frac{1}{2}, \bar{z}; 0, \frac{1}{2}, \frac{1}{2} + z; 0, \frac{1}{2}, \frac{1}{2} - z;$ $\frac{1}{2}, 0, z; \frac{1}{2}, 0, \bar{z}; \frac{1}{2}, 0, \frac{1}{2} + z; \frac{1}{2}, 0, \frac{1}{2} - z.$	$hkl: h+k=2n, l=2n$
4	g	mm	$x, \bar{x}, 0; \bar{x}, x, 0; \frac{1}{2} + x, \frac{1}{2} + x, \frac{1}{2}; \frac{1}{2} - x, \frac{1}{2} - x, \frac{1}{2}.$	
4	f	mm	$x, x, 0; \bar{x}, \bar{x}, 0; \frac{1}{2} + x, \frac{1}{2} - x, \frac{1}{2}; \frac{1}{2} - x, \frac{1}{2} + x, \frac{1}{2}.$	
4	e	mm	$0, 0, z; 0, 0, \bar{z}; \frac{1}{2}, \frac{1}{2}, \frac{1}{2} + z; \frac{1}{2}, \frac{1}{2}, \frac{1}{2} - z.$	
4	d	$\bar{4}$	$0, \frac{1}{2}, \frac{1}{4}; \frac{1}{2}, 0, \frac{1}{4}; 0, \frac{1}{2}, \frac{3}{4}; \frac{1}{2}, 0, \frac{3}{4}.$	
4	c	$2/m$	$0, \frac{1}{2}, 0; \frac{1}{2}, 0, 0; 0, \frac{1}{2}, \frac{1}{2}; \frac{1}{2}, 0, \frac{1}{2}.$	
2	b	mmm	$0, 0, \frac{1}{2}; \frac{1}{2}, \frac{1}{2}, 0.$	
2	a	mmm	$0, 0, 0; \frac{1}{2}, \frac{1}{2}, \frac{1}{2}.$	






Quantum mechanics at work





thanks to Erich Wimmer




TiC electron density



- NaCl structure (100) plane
- Valence electrons only
- plot in 2 dimensions
- Shows
 - *charge distribution*
 - *covalent bonding*
 - between the Ti-3d and C-2p electrons
 - e_g/t_{2g} symmetry






TiC, three valence states at Δ



Energy bands











Ti-4s

Ti-3d

C-2p

C-2s



Δ_1 423mRyd

Δ_2' 620mRyd

Δ_5 636mRyd


$C_p-Ti_d \sigma$

$Ti_d-Ti_d \sigma$

$C_p-Ti_d \pi$

(100) plane

P.Blaha, K.Schwarz,
Int.J.Quantum Chem. 23, 1535 (1983)




TiC, energy bands




spaghetti

irred.rep.


character bands



P.Blaho, K.Schwarz,
Int.J.Quantum Chem. 23, 1535 (1983)






TiC, bonding and antibonding states



Ti-3d

O-2p



weight:



P.Blaha, K.Schwarz,
Int.J.Quantum Chem. 23, 1535 (1983)





Bonding and antibonding state at Δ_1




antibonding
 $C_p-Ti_d \sigma$



bonding
 $C_p-Ti_d \sigma$





TiC, TiN, TiO





Figure 3. Valence charge densities in the (100) plane. Contour intervals $0.1e\text{\AA}^{-3}$ (numbers are in these units), cutoff at $1.7e\text{\AA}^{-3}$.

TiC

TiN

TiO


Rigid band model: **limitations**

Electron density ρ : **decomposition**

$$1 = q_{out} + \sum_t \sum_l q_{tl}$$

unit cell interstitial atom t $l=s, p, d, \dots$

P.Blaha, K.Schwarz,
Int.J.Quantum Chem. 23, 1535 (1983)






TiC, TiN, TiO



Atomic form factors for Ti and C

Experimental difference electron density



Paired reflections

$$s = |\vec{S}| \approx \frac{\sin \vartheta}{\lambda}$$

\vec{S}			$h^2+k^2+l^2$
h	k	l	
10	2	2	108
6	6	6	108

$$F(\vec{S}) = F(S) \quad \text{spheric. symm. density}$$

$$\left. \begin{aligned} F(\vec{S}_1) &\neq F(\vec{S}_2) \\ \text{with } |\vec{S}_1| &= |\vec{S}_2| \end{aligned} \right\} \text{non spherical}$$



Vienna, city of music and the **Wien2k** code

